

Seed Potatoes and Tissue-Culture Plantlets: Health Evidence, Establishment and Procurement Validation

Connect material identity and health testing with the purchase, delivery, receipt and establishment of in-vitro plantlets, acclimatized plants, minitubers and field-grown seed. Interpret generation labels, negative tests and field performance within their evidence limits.

Junsheng

Version 1.0 | 2026-10-16

For general technical discussion and procurement reference. This document does not replace product specifications, batch testing or validation in the actual process. Product requirements and supply commitments follow the agreed technical documents and contract.

No unauthorized reproduction, distribution, republication, adaptation or commercial republication of copyright-protected original content. Downloading, reading and internal procurement evaluation are permitted. Statutory exceptions are preserved. Rights in quoted third-party material remain with its owners.

Contents

01 Identify the material and the next production stage	3
02 Variety, generation and lot are different identity records	4
03 Distinguish virus elimination, aseptic culture and a negative test	5
04 Why no detection does not prove an entirely pest-free lot	6
05 Receive and acclimatize plantlets as an environmental transition	7
06 Manage physiological condition and health during storage	8
07 Record quality and the timing of handover	8
08 Separate material effects from field conditions	10
09 Compare cost per usable unit and establishment outcome	11
10 Close the record from source to delivery and feedback	11
11 Worked establishment example and the next purchasing cycle	12

01 Identify the material and the next production stage

The first purchasing decision is whether the material matches the next production stage.

In-vitro plantlets, acclimatized plants, minitubers produced under protected conditions and field-grown seed tubers require different receiving arrangements and management. Calling all of them “seed plants” can hide hardening, isolation or dormancy-management requirements.

Buyers of in-vitro plantlets need the capacity to receive, inspect, acclimatize and multiply them. For acclimatized plants, confirm hardening stage, root condition and transport format. For tubers, also consider size distribution, sprout condition, storage history and intended planting date. A low purchase price per unit does not establish a low cost per usable plant or successfully planted area.

Junsheng produces seed potatoes and tissue-culture plantlets. The variety, material form, multiplication stage, lot and delivery conditions are confirmed in the relevant supply documents. Production ownership does not establish freedom from every pathogen. A variety name does not guarantee yield, processing performance or resistance in every environment.

Express the requirement as a usable quantity of a defined material, needed at a specified place and date, under identified protected or field conditions, for a particular multiplication or commercial-crop stage. That statement guides material selection, delivery scheduling and acceptance more effectively than a general request for “high-yield virus-free seed”.

Material	Next operation	Specify in the order
In-vitro plantlets	Receipt, inspection, acclimatization	Container, condition, lot, health report
Acclimatized plants	Holding and establishment	Hardening stage, roots, tray or substrate
Minitubers	Storage, sprouting and multiplication	Production system, size, date
Field seed tubers	Planting and field management	Scheme, generation, size, health and storage history

02 Variety, generation and lot are different identity records

Variety identifies the genetic material. Generation or class locates it within a particular multiplication or certification scheme. Lot identity connects a delivery to production, testing and handover records. One variety can occur at several generations; one generation can contain lots with different health and storage histories.

Names used by countries, certification authorities and production systems are not automatically interchangeable. Before mapping terms such as “G0” or a domestic class, examine the scheme, starting material, multiplication history and documented definition. The UNECE seed-potato framework addresses varietal identity, genealogy, health, external quality and labelling, but does not automatically replace destination requirements.[P1]

When checking a delivery lot, buyers can use a relationship record connecting starting-material ID, variety, stage, production location and date, split or merged lots, test-sample ID and shipping label. Retaining one old sample number after combining material can leave the customer unable to determine what the report covers. Repacking without changing material still requires a link to the original lot.

Appearance alone does not establish varietal identity, and a health test does not evaluate every genetic attribute. For identity disputes or specialized processing uses, select the

documents and methods appropriate to that question. Do not extend a report to attributes it did not measure.

03 Distinguish virus elimination, aseptic culture and a negative test

Virus elimination describes treatment and selection directed at specified viral targets. Aseptic culture describes microbial control in the culture environment. A negative test is a result for specified targets, specimens and methods. These concepts are connected but not interchangeable. A visibly clean vessel does not demonstrate absence of all viruses, and treated material still requires testing and maintained protection.

In-vitro conservation and multiplication must preserve identity as well as health. CIP's description of its collection highlights both.[P5] The purchasing task is not to assume that a supplier follows another institution's system, but to connect delivered material with its actual testing and maintenance records.

Read the report for the target organisms, material type, sampling date and locations, sample or pool numbers, method and interpretation. The method must suit the pathogen and specimen matrix. EPPO organizes diagnostic protocols by target and method context.[P4] A general “virus negative” heading without a target list does not define the tested health scope.

Multiplication, transfers and exposure after sampling affect what a result can support, including changes before the report is issued. A single test on starting plantlets cannot automatically cover descendants produced through multiple generations or environments. Specify whether the purchasing requirement concerns starting material, the delivery lot or a defined production stage.

04 Why no detection does not prove an entirely pest-free lot

Detection in a sampled lot depends on prevalence, spatial distribution, sampling and test capability. If affected material is clustered in one vessel or production area, repeated testing at other locations may fail to find it. Pooling can dilute the target and must follow a validated method and pool size.

The following is a probability illustration, not a phytosanitary sampling plan. Assume a large population, independent units, random sampling, prevalence p , probability s of detecting an infected sampled unit, and no false positives for the calculation. The probability of at least one detection among n units is $1 - (1 - p \times s)^n$. Do not transfer the result when those assumptions do not hold.

At a hypothetical prevalence of 1% and perfect detection, testing 100 units provides approximately 63.4% probability of at least one detection; testing 300 provides approximately 95.1%. Lower sensitivity, uneven sampling or clustering changes performance. All-negative observations do not prove that the lot contains no infection. These detection probabilities are not probabilities that a lot is infection-free after negative results, nor are they certification thresholds.

ISPM 31 addresses sampling methodologies for consignment inspection or testing by national plant protection organizations; its scope excludes field-survey sampling.[P3] Apply the requirements of the competent authority and relevant scheme for phytosanitary certification. A buyer's additional incoming-quality check does not replace that official process.

05 Receive and acclimatize plantlets as an environmental transition

Transfer from an in-vitro vessel to substrate or another production system changes humidity, light, air movement, water supply and the root environment simultaneously. Acclimatization must fit variety, development stage, root condition, season and facilities. Cultivar-specific research evaluates acclimatization, later performance and fidelity separately; a reported survival result for one cultivar is not a general promise for supplied plantlets.[P6]

At receipt, inspect container integrity, labels, count, medium or substrate condition, and visible leaf, stem and root abnormalities. Record transport damage, suspected contamination, dehydration and uneven growth separately. Avoid mixing unchecked material into accepted lots. Photographs should show the package, label and representative abnormalities with date information.

After establishment, record more than survival. At predefined observation times distinguish living plants, plants resuming growth and plants fit for further multiplication. A green plant without new growth is not equivalent to one with established roots and active development. State whether the denominator is the quantity delivered, actually planted or still alive when observed.

Validate a receiving and acclimatization route on representative material before expanding. Record environment and operational changes. Do not adopt one research study's humidity or temperature programme as a universal protocol, or attribute every abnormality to a single action such as watering, shading or nutrition.

06 Manage physiological condition and health during storage

Equal harvest dates do not establish equal physiological condition. Variety, maturity, temperature history, sprout condition and handling damage influence readiness for planting. In addition to quantity and size, specify the interval between expected receipt and planting and the buyer's ventilation, temperature-management and lot-segregation capacity.

Cold tubers entering warm humid surroundings can develop condensation. Limited airflow, local wetness and wounds complicate quality preservation. AHDB guidance emphasizes identification, ventilation and condensation management.[P7][P8] The receiving team can use these principles to check lot labels, ventilation paths and locations susceptible to condensation.

Distinguish room-air readings, packaging surroundings and representative tuber positions. A normal logger reading represents its location; it does not prove uniform conditions throughout the load. Inspect different locations for localized moisture, odour, rot, abrasion, sprout damage and size distribution, retaining the connection between abnormalities and package position.

Agree subsequent handling according to material condition, variety and planting schedule. Do not transfer sprout-suppression treatments intended for consumption potatoes directly to seed. Select storage temperature and sprouting duration for variety, dormancy and physiological condition, storage history and planting date. An unsuitable treatment can undermine vigour or disrupt the planting programme.

07 Record quality and the timing of handover

Connect four periods in the receiving record: before dispatch, transport, arrival and handling after receipt. A photograph taken several days later rarely establishes when a problem began. Lot identity, seals, package position, transport records, arrival inspection and the receiving environment together support investigation.

Visual quality, health testing and establishment performance answer different questions. Visually normal material may still require target-specific testing. Material negative within the tested scope can nevertheless perform poorly after dehydration, sprout damage or unsuitable receiving conditions. A single “quality satisfactory” tick discards information needed later.

Identify and segregate abnormal material, retain original labels and evidence, and contact the supplier through the agreed procedure. Refer health or phytosanitary concerns to appropriate professionals. Do not mix suspect material into production while waiting to resolve its status. Use, retesting, replacement or return depends on the problem, applicable requirements and contract.

A sampled defect rate initially describes the inspected sample. Inference to the lot requires representative sampling, a suitable design and consideration of sampling uncertainty. A statement such as “2% damaged” needs an inspection count, defect definition, counting rule and sampling locations. Several defects may occur on one tuber; adding their individual percentages does not necessarily give the total proportion of unacceptable tubers.

Check	Record	Does not replace
Identity/count	Labels, lot, form, quantity	All varietal-identity evidence
Condition	Defined defects, positions, photos	Pathogen testing
Health documents	Targets, sampling, methods, dates	Post-receipt management
Transport	Times, logger positions, packaging	History of every unit
Disputed material	Segregation, samples, retest records	Applicable official or contractual procedure

08 Separate material effects from field conditions

Good performance in one field does not by itself prove that replacing planting material increases yield. Soil, topography, irrigation, pest pressure, planting date, nutrition and harvest timing can all influence results. Keep these conditions comparable where possible, use independently replicated randomized plots, and record weather or management deviations. AHDB guidance for determining varietal nitrogen groups also emphasizes seed comparability and randomized replication.[P9] When comparing lots of the same variety, control factors such as seed size and physiological condition. Comparisons across varieties also contain genetic differences and cannot attribute every effect to health status.

The experimental unit is an independently allocated plot, not every plant within one plot. Establish rules for border plants, gaps and replacements before observing the outcome. Follow emergence, establishment, stand retention, health observations, yield and size distribution over time. Processing applications also require appropriate measurement of dry matter, processing colour or other agreed endpoints.

Use actual harvested area as the yield denominator. Report total yield and yield within the required marketable or usable categories. Do not select only the most favourable size band or ignore unequal harvest dates. For seed multiplication, health- and size-acceptable output may matter more than total fresh mass.

One site-season supports a conclusion under those conditions, not a universal regional yield improvement. Broader claims require trials relevant to intended locations, seasons and management. When performance is poor, locate the divergence in emergence, stand formation, health or tuber distribution rather than assigning every failure to seed degeneration.

09 Compare cost per usable unit and establishment outcome

Quotations for in-vitro plants, acclimatized plants and tubers use different units. A direct comparison of prices per plant or tonne can mislead. Define the final usable unit: a plant ready for the next stage, an acceptable seed tuber or successfully established area. Include purchase, transport, receipt, acclimatization or storage and attributable losses within one consistent boundary.

Illustrative calculation: 10,000 delivered plants cost 12,000 currency units including receipt-related costs. After a predefined recovery period, 9,000 meet the next-stage criterion. Cost per usable plant within that boundary is 1.33. Another option costing 13,000 with 9,800 usable plants is also approximately 1.33. A higher quoted unit price does not necessarily create a higher usable-unit cost. These are hypothetical inputs, not Junsheng survival rates or customer results.

Economic comparison begins only after identity, health and suitability requirements are met. It cannot justify relaxing them. Time to production, facility occupancy and uneven growth caused by replacement planting can also carry cost; use recorded evidence rather than assuming unmeasured items are zero.

For successive multiplication stages, distinguish increasing quantity from maintaining acceptable health and quality. A high multiplication factor accompanied by more rejects, longer cycles or fewer units in the target grade may not reduce final usable-unit cost.

10 Close the record from source to delivery and feedback

A verifiable purchasing file links material identity, production origin, health evidence, transport handover and subsequent use to the same lot. Applicable Junsheng records concerning source material, incoming checks, process control and finished-product verification are linked through blockchain records for review. That linkage does not independently establish the truth of the source data or replace testing and official certification.

For international projects, check requirements for the material form, origin and destination. ISPM 33 concerns potato micropropagative material and minitubers intended for international trade; it explicitly excludes field-grown material and potatoes for consumption or processing.[P2] It is therefore not a universal certification for every seed-potato delivery. Import conditions, phytosanitary certificates and permit arrangements depend on the relevant authorities.

Before ordering, establish four connections: the ordered material matches the documents; the tested specimens match the production stage or lot claimed; delivery timing matches receiving capacity; and acceptance methods match the agreed response procedure. If a report covers starting material only, document the multiplication relationship rather than presenting its date as a dispatch-date test.

Return meaningful production feedback with its conditions: lot, location, planting date, environment, management, emergence records, abnormality photographs and harvest endpoints. Keeping results with these conditions enables the next purchase to distinguish material effects, operational improvements and seasonal variation.

11 Worked establishment example and the next purchasing cycle

“High survival” can use very different denominators. In this fictional receiving example, 10,000 plantlets arrive and 9,800 pass visual receipt checks. The buyer plants 9,600 and retains 200 as unplanted reserves. At a predefined observation date, 9,000 satisfy morphological and growth criteria for further multiplication. The qualifying proportion is $9,000 \div 9,600 = 93.75\%$ of planted units and 90% of delivered units. Both can be reported with definitions; neither establishes a negative result for every pathogen.

The record distinguishes 200 units not accepted on arrival, 200 not planted, and 600 planted units not reaching the observation criterion. These outcomes represent different management questions and cannot all be called transport loss. Recording abnormalities by vessel or tray allows investigation of clustering in packaging, receiving groups or facility locations.

This makes the next purchasing improvement specific. Packaging-associated defects call for review of loading and transport. A problem concentrated during an acclimatization transition calls for a small controlled comparison. Abnormalities shared across locations call for examination of common substrate, water or management changes and professional judgment on targeted health testing. This is an investigation sequence, not a pathogen diagnosis from symptoms alone.

The completed file should identify the material lot, arrival time, sampling procedure, establishment date, evaluation criteria, usable output and changes needing validation next time. This record supports dependable supply more effectively than an undefined percentage.

Stage	Illustrative count	Interpretation
Delivered	10,000	Delivery denominator
Visually accepted	9,800	Not universal pathogen clearance
Planted	9,600	200 reserves excluded from planting denominator
Meeting criterion	9,000	Specify observation date and criterion
Planted, not qualifying	600	Classify and locate; do not assume supplier cause

References

- P1. UNECE. Seed Potatoes: Standard S-1 and certification resources.
<https://unece.org/trade/wp7/SeedPotatoes-Standards>
- P2. IPPC. ISPM 33: Pest free potato micropropagative material and minitubers for international trade.
<https://www.ippc.int/en/publications/pest-free-potato-solanum-spp-micropropagative-material-and-minitubers-international/>
- P3. IPPC. ISPM 31: Methodologies for sampling of consignments.
<https://www.ippc.int/en/publications/588/>
- P4. EPPO. PM 7 diagnostic protocols for regulated pests.
<https://gd.eppo.int/standards/pm7/>
- P5. CIP Genebank. In Vitro Collection.
https://cipotato.org/genebankcip/process/active_collection/
- P6. A mono-phasic protocol for micropropagation of potato cv. Cooch Behar local, its acclimatization, on-field evaluation, and fidelity analysis.
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11751352/>
- P7. AHDB. Seed handling & storage.
<https://potatoes.ahdb.org.uk/knowledge-library/seed-handling-storage>
- P8. AHDB. Storage practices to minimise the risk of soft rot and blackleg in seed potatoes.
<https://horticulture.ahdb.org.uk/knowledge-library/storage-practices-to-minimise-the-risk-of-soft-rot-and-blackleg-in-seed-potatoes>
- P9. AHDB. In-field method for assigning nitrogen group (determinacy) in potatoes.
<https://potatoes.ahdb.org.uk/knowledge-library/in-field-method-for-assigning-nitrogen-group-determinacy-in-potatoes>